

Two-particle correlations in azimuthal angle and pseudorapidity in inelastic p+p interactions at the CERN Super Proton Synchrotron

A. Aduszkiewicz¹⁷, Y. Ali^{15,32}, E. Andronov²³, T. Antić³, N. Antoniou⁸, B. Baatar²¹, F. Bay²⁵,
A. Blondel²⁷, M. Bogomilov², A. Brandin²², A. Bravar²⁷, J. Brzychczyk¹⁵, S.A. Bunyatov²¹,
O. Busygina²⁰, P. Christakoglou⁸, M. Ćirković²⁴, T. Czopowicz¹⁹, A. Damyanova²⁷, N. Davis¹³,
H. Dembinski⁵, M. Deveau⁷, F. Diakonov⁸, S. Di Luise²⁵, W. Dominik¹⁷, J. Dumarchez⁴,
R. Engel⁵, A. Ereditato²⁶, G.A. Feofilov²³, Z. Fodor^{9,18}, A. Garibov¹, M. Gaździcki^{7,12},
M. Golubeva²⁰, K. Grebieszko¹⁹, A. Grzeszczuk¹⁶, F. Guber²⁰, A. Haesler²⁷, T. Hasegawa¹⁰,
A.E. Hervé⁵, M. Hierholzer²⁶, J. Hlyen²⁸, S. Igoikin²³, A. Ivashkin²⁰, S.R. Johnson³⁰, K. Kadja³,
A. Kapoyannis⁸, E. Kaptur¹⁶, M. Kielbowicz¹³, J. Kisiel¹⁶, N. Knezević²⁴, T. Kobayashi¹⁰,
V.I. Kolesnikov²¹, D. Kolev², V.P. Kondratiev²³, A. Korzenev²⁷, V. Kovalenko²³, K. Kowalik¹⁴,
S. Kowalski¹⁶, M. Koziel⁷, A. Krasnoperov²¹, M. Kuich¹⁷, A. Kurepin²⁰, D. Larsen¹⁵, A. László⁹,
M. Lewicki¹⁸, B. Lundberg²⁸, V.V. Lyubushkin²¹, M. Maćkowiak-Pawłowska¹⁹, B. Maksia¹⁹,
A.I. Malakhov²¹, D. Manić²⁴, A. Marchionni²⁸, A. Marcinek^{15,18}, A.D. Marino³⁰, K. Marton⁹,
H.-J. Mathes⁵, T. Matulewicz¹⁷, V. Matveev²¹, G.L. Melkumov²¹, A. Merzlaya²³, B. Messerly³¹,
G.B. Mills²⁹, S. Morozov^{20,22}, S. Mrówczyński¹², Y. Nagai³⁰, T. Nakadaira¹⁰, M. Naskręć¹⁸,
M. Nirkko²⁶, K. Nishikawa¹⁰, V. Ozvenchuk¹³, A.D. Panagiotou⁸, V. Paolone³¹, M. Pavin^{4,3},
O. Petukhov^{20,22}, C. Pistillo²⁶, R. Płaneta¹⁵, B.A. Popov^{21,4}, M. Posiadła¹⁷, S. Puławski¹⁶,
J. Puzović²⁴, R. Rameika²⁸, W. Rauch⁶, M. Ravonel²⁷, A. Redij²⁶, R. Renfordt⁷, E. Richter-Was¹⁵,
A. Robert⁴, D. Röhrich¹¹, E. Rondio¹⁴, M. Roth⁵, A. Rubbia²⁵, B.T. Rumberger³⁰, A. Rustamov^{1,7},
M. Rybczynski¹², A. Rybicki¹³, A. Sadovsky²⁰, K. Sakashita¹⁰, R. Sarnecki¹⁹, K. Schmidt¹⁶,
T. Sekiguchi¹⁰, I. Selyuzhenkov²², A. Seryakov²³, P. Seyboth¹², D. Sgalaberna²⁵, M. Shibata¹⁰,
M. Słodkowski¹⁹, P. Staszal¹⁵, G. Stefanek¹², J. Stepaniak¹⁴, H. Ströbele⁷, T. Šuš³, M. Szuba⁵,
M. Tada¹⁰, A. Taranenko²², A. Tefelska¹⁹, D. Tefelski¹⁹, V. Tereshchenko²¹, R. Tsenov²,
L. Turko¹⁸, R. Ulrich⁵, M. Unger⁵, M. Vassiliou⁸, D. Veberič⁵, V.V. Vechernin²³, G. Vesztegombi⁹,
L. Vinogradov²³, M. Walewski¹⁷, A. Wickremasinghe³¹, A. Wilczek¹⁶, Z. Włodarczyk¹²,
A. Wojtaszek-Szwarc¹², O. Wyszynski¹⁵, L. Zambelli^{4,10}, E.D. Zimmerman³⁰, R. Zwaska²⁸

¹National Nuclear Research Center, Baku, Azerbaijan

²Faculty of Physics, University of Sofia, Sofia, Bulgaria

³Ruder Bošković Institute, Zagreb, Croatia

⁴LPNHE, University of Paris VI and VII, Paris, France

⁵Karlsruhe Institute of Technology, Karlsruhe, Germany

⁶Fachhochschule Frankfurt, Frankfurt, Germany

⁷University of Frankfurt, Frankfurt, Germany

⁸University of Athens, Athens, Greece

⁹Wigner Research Centre for Physics of the Hungarian Academy of Sciences, Budapest, Hungary

¹⁰Institute for Particle and Nuclear Studies, Tsukuba, Japan

¹¹University of Bergen, Bergen, Norway

¹²Jan Kochanowski University in Kielce, Poland

¹³H. Niewodniczański Institute of Nuclear Physics of the Polish Academy of Sciences, Kraków, Poland

¹⁴National Centre for Nuclear Research, Warsaw, Poland

¹⁵Jagiellonian University, Cracow, Poland

¹⁶University of Silesia, Katowice, Poland

¹⁷University of Warsaw, Warsaw, Poland

¹⁸University of Wrocław, Wrocław, Poland

¹⁹Warsaw University of Technology, Warsaw, Poland

²⁰Institute for Nuclear Research, Moscow, Russia

²¹Joint Institute for Nuclear Research, Dubna, Russia

²²National Research Nuclear University (Moscow Engineering Physics Institute), Moscow, Russia

²³St. Petersburg State University, St. Petersburg, Russia

²⁴University of Belgrade, Belgrade, Serbia

²⁵ETH Zürich, Zürich, Switzerland

²⁶University of Bern, Bern, Switzerland

²⁷University of Geneva, Geneva, Switzerland

²⁸Fermilab, Batavia, USA

²⁹Los Alamos National Laboratory, Los Alamos, USA

³⁰University of Colorado, Boulder, USA

³¹University of Pittsburgh, Pittsburgh, USA

³²Present address: COMSATS Institute of Information Technology, Islamabad, Pakistan

Received: date / Accepted: date

Abstract Results on two-particle $\Delta\eta\Delta\phi$ correlations in inelastic p+p interactions at 20, 31, 40, 80, and 158 GeV/c are presented. The measurements were performed using the large acceptance NA61/SHINE hadron spectrometer at the CERN Super Proton Synchrotron. The data show structures which can be attributed mainly to effects of resonance decays, momentum conservation, and quantum statistics. The results are compared with the EPOS and UrQMD models.

Keywords two-particle correlations · azimuthal angle · pseudorapidity · resonance decays · conservation laws · flux-tube fragmentation

PACS 25.75.-q, 25.75.Gz

1 Introduction and motivation

This paper presents experimental results on two-particle correlations in pseudorapidity and azimuthal angle of charged particles produced in inelastic p+p interactions at 20, 31, 40, 80, and 158 GeV/c. The measurements were performed by the multi-purpose NA61/SHINE [1, 2] experiment at the CERN Super Proton Synchrotron (SPS). They are part of the strong interactions programme devoted to the study of the properties of the onset of deconfinement and search for the critical point of strongly interacting matter. Within this program a two dimensional scan in collision energy and size of colliding nuclei is in progress. Data on p+p, Be+Be, and Ar+Sc collisions were already recorded and data on p+Pb and Xe+La collisions will be registered within the coming years. The expected signal of a critical point is a non-monotonic dependence of various fluctuation measures in such a scan, for a recent review see Ref. [3].

Apart from looking for QGP signatures, it is of interest to study specific physical phenomena that happen during and after the collision. The two-particle correlation analysis in pseudorapidity (η) and azimuthal angle (ϕ) allows to disentangle different correlation sources which can be directly connected with phenomena like jets, collective flow, resonance decays, quantum statistics effects, conservation laws, etc.

Measurements of two-particle correlations in pseudorapidity and azimuthal angle were first published by the ACM collaboration at the Intersecting Storage Rings (ISR) [4]. Two- and three-body decays of resonances (η , ρ^0 , ω) were found to provide the dominant contributions. Two structures were discovered: an enhancement near $\Delta\phi = \pi$ (away-side) explained by the two-body decay scenario and another enhancement at $\Delta\phi \approx 0$ together with an azimuthal ridge (centered at $\Delta\eta \approx 0$) consistent with three-body decays¹. These features were confirmed at the higher collision energies of RHIC by the PHOBOS [5] collaboration.

At RHIC and the LHC parton scattering processes become important. In addition to high transverse momentum jets, studies of $\Delta\eta\Delta\phi$ correlations in p+p interactions as well as in collisions

of heavy nuclei [6–9] found prominent structures explained as arising from the production of minijets, producing a large correlation peak at small opening angles ($\Delta\eta, \Delta\phi$) \approx (0, 0) and a broad structure along $\Delta\eta$ at $\Delta\phi \approx \pi$ (also referred to as away-side ridge). At SPS energies, however, contributions from hard and semi-hard scattering processes are expected to be much smaller.

Collective effects also provide significant contributions to two-particle correlations. They appear as a modulation in $\Delta\phi$ (see e.g. Ref. [10]). The effect of flow is most visible in heavy-ion collisions but also appears in high-multiplicity p+p [11, 12] and p+Pb [13] interactions. At SPS energies sufficiently high particle multiplicities cannot be produced in p+p collisions. Thus the effect will not be discussed in more detail in this paper.

In this paper the pseudorapidity variable η is calculated as $\eta = -\log \tan(\Theta/2)$ where $\tan(\Theta/2) = p_T/p_L$ with p_T the transverse (x, y) and p_L the longitudinal (z) component of the particle momentum in the collision centre of mass system. The azimuthal angle ϕ is the angle between the transverse momentum vector and the horizontal (x) axis.

The paper is organized as follows. The correlation function is defined in Sec. 2. The experimental setup is presented in Sec. 3. Data processing and simulation are described in Sec. 4. Data selection and analysis are discussed in Sec. 5. Results are presented in Sec. 6 and compared to model calculations in Sec. 7. A summary in Sec. 8 closes the paper.

2 Two-particle correlations in pseudorapidity and azimuthal angle

Correlations studied in this paper were calculated as a function of the difference in pseudorapidity (η) and azimuthal angle (ϕ) between two particles produced in the same event:

$$\Delta\eta = |\eta_1 - \eta_2|, \quad \Delta\phi = |\phi_1 - \phi_2|. \quad (1)$$

The correlation function $C(\Delta\eta, \Delta\phi)$ is defined and calculated as:

$$C(\Delta\eta, \Delta\phi) = \frac{N_{\text{mixed}}^{\text{pairs}} D(\Delta\eta, \Delta\phi)}{N_{\text{data}}^{\text{pairs}} M(\Delta\eta, \Delta\phi)}, \quad (2)$$

where

$$D(\Delta\eta, \Delta\phi) = \frac{d^2 N_{\text{data}}}{d\Delta\eta d\Delta\phi}, \quad M(\Delta\eta, \Delta\phi) = \frac{d^2 N_{\text{mixed}}}{d\Delta\eta d\Delta\phi}$$

are the distributions of particle pairs from data and mixed events (representing uncorrelated background), respectively. The mixed events were constructed using the mixing procedure described in Sec. 5.1. Distributions $D(\Delta\eta, \Delta\phi)$ and $M(\Delta\eta, \Delta\phi)$ were obtained by accumulating the number of pairs in intervals of $\Delta\eta$ and $\Delta\phi$. For the calculation of $C(\Delta\eta, \Delta\phi)$ both distributions were normalised to the number of pairs ($N_{\text{data}}^{\text{pairs}}$, $N_{\text{mixed}}^{\text{pairs}}$) in the given distribution. The measurements are restricted to $0 \leq \Delta\eta \leq 3$ and $0 \leq \Delta\phi < \pi$. Finally the correlation functions were mirrored around

¹ $\Delta\eta$ and $\Delta\phi$ definitions are in Eq. 1

the point $(\Delta\eta, \Delta\phi) = (0, 0)$ and plotted in the range $-\frac{\pi}{2} < \Delta\phi < \frac{3\pi}{2}$.

In this paper the correlation function $C(\Delta\eta, \Delta\phi)$ was calculated for charged primary hadrons produced in strong and electromagnetic processes with transverse momentum $p_T < 1.5$ GeV/c in the NA61/SHINE acceptance (for details see [14]).

While CMS [11], ATLAS [15] and ALICE [16] used equivalent definitions, other quantities have also been used to measure two-particle correlations. The ACM collaboration [4] at the ISR studied

$$C^H(\Delta\eta, \Delta\phi) = \langle (n-1)(\rho_n^H(\eta_1, \phi_1, \eta_2, \phi_2) - \rho_n^I(\eta_1, \phi_1)\rho_n^I(\eta_2, \phi_2)) \rangle,$$

where n is the multiplicity in the event, ρ_n^H the pair density, ρ_n^I the inclusive density and the averaging runs over the events.

PHOBOS [5] employed the observable

$$R(\Delta\eta, \Delta\phi) = \langle (n-1)(\rho_n^H(\Delta\eta, \Delta\phi)/\rho_n^{\text{mixed}}(\Delta\eta, \Delta\phi) - 1) \rangle$$

with ρ_n^H and ρ_n^{mixed} the pair density in data and mixed events, respectively, both normalised to unity.

The STAR collaboration [6] used the quantity

$$\Delta\rho/\sqrt{\rho_{\text{mixed}}} = (\rho(\Delta\eta, \Delta\phi) - \rho_{\text{mixed}}(\Delta\eta, \Delta\phi))/\sqrt{\rho_{\text{mixed}}(\Delta\eta, \Delta\phi)},$$

where ρ and ρ_{mixed} are the normalised densities of data and mixed pairs respectively.

3 Experimental setup, beams, target, triggers

The NA61/SHINE experiment uses a large acceptance hadron spectrometer located in the CERN North Area [2]. The schematic layout of the NA61/SHINE detector is shown in Fig. 1.

A set of scintillation and Cherenkov counters as well as beam position detectors (BPDs) upstream of the spectrometer provide timing reference, identification and position measurements of incoming beam particles. The trigger scintillator counter S4 placed downstream of the target is used to select events with collisions in the target area by the absence of a charged particle hit.

Secondary beams of positively charged hadrons at 20, 31, 40, 80, and 158 GeV/c are produced from 400 GeV/c protons extracted from the SPS accelerator. Particles of the secondary hadron beam are identified by two Cherenkov counters, a CEDAR [17] (either CEDAR-W or CEDAR-N) and a threshold counter (THC). The CEDAR counter, using a coincidence of six out of the eight photo-multipliers placed radially along the Cherenkov ring, provides positive identification of protons, while the THC, operated at pressure lower than the proton threshold, is used in anti-coincidence in the trigger logic. Due to their limited range of operation two different CEDAR counters were used, namely for beams at 20, 31, and 40 GeV/c the CEDAR-W counter and for beams at 80 and 158 GeV/c the CEDAR-N counter. The threshold counter was used for the lower three beam momenta. A selection based on signals from the Cherenkov counters allowed to identify

beam protons with a purity of about 99%. A consistent value for the purity was found by bending the beam into the TPCs with the full magnetic field and using identification based on its specific ionization energy loss dE/dx [18].

The main tracking devices of the spectrometer are four large volume Time Projection Chambers (TPCs). Two of them, the vertex TPCs (VTPC-1 and VTPC-2), are located in the magnetic fields of two super-conducting dipole magnets with a maximum combined bending power of 9 Tm which corresponds to about 1.5 T and 1.1 T fields in the upstream and downstream magnets, respectively. In order to optimize the acceptance of the detector, the field in both magnets was adjusted proportionally to the beam momentum.

Two large *main* TPCs (MTPC-L and MTPC-R) are positioned downstream of the magnets symmetrically to the beam line. The fifth small TPC (GAP-TPC) is placed between VTPC-1 and VTPC-2 directly on the beam line. It closes the gap between the beam axis and the sensitive volumes of the other TPCs. The TPCs are filled with Ar and CO₂ gas mixtures. Particle identification in the TPCs is based on measurements of the specific energy loss (dE/dx) in the chamber gas and is augmented by mass measurements using Time-of-Flight (ToF) detectors.

The p+p data, which is the topic of this paper, was taken by colliding the proton beam with a liquid hydrogen target (LHT), a 20 cm long cylinder positioned about 80 cm upstream of VTPC-1.

4 Data processing, simulation and detector performance

Detector parameters were optimised by a data-based calibration procedure which also took into account their time dependencies. Small adjustments were determined in consecutive steps for:

- (i) detector geometry, TPC drift velocities and distortions due to the magnetic field inhomogeneities in the corners of the VTPCs,
- (ii) magnetic field setting,
- (iii) specific energy loss measurements.

Each step involved reconstruction of the data required to optimise a given set of calibration constants and time dependent corrections followed by verification procedures. Details of the procedure and quality assessment are presented in Ref. [19]. The resulting performance in the measurements of quantities relevant for this paper is discussed below.

The main steps of the data reconstruction procedure are:

- (i) cluster finding in the TPC raw data, calculation of the cluster centre-of-gravity and total charge,
- (ii) reconstruction of local track segments in each TPC separately,
- (iii) matching of track segments into global tracks,
- (iv) track fitting through the magnetic field and determination of track parameters at the first measured TPC cluster,




Fig. 1 (Color online) The schematic layout of the NA61/SHINE experiment at the CERN SPS (horizontal cut, not to scale).

- (v) determination of the interaction vertex using the beam trajectory (x and y coordinates) fitted in the BPDs and the trajectories of tracks reconstructed in the TPCs (z coordinate),
- (vi) refitting the particle trajectory using the interaction vertex as an additional point and determining the particle momentum at the interaction vertex.

A simulation of the NA61/SHINE detector response is used to correct the reconstructed data. For this purpose inelastic p+p collisions generated with the EPOS 1.99 [20] model were used to obtain the corrections for reconstruction inefficiency and trigger-bias of the NA61/SHINE detector and obtain final corrected results.

The simulation consists of the following steps (see Ref. [21] for more details):

- (i) generation of inelastic p+p interactions using the EPOS model,
- (ii) propagation of outgoing particles through the detector material using the GEANT 3.21 package [22] which takes into account the magnetic field as well as relevant physics processes, such as particle interactions and decays,
- (iii) simulation of the detector response using dedicated NA61/SHINE packages which introduce distortions corresponding to all corrections applied to the real data,
- (iv) simulation of the interaction trigger selection by checking whether a charged particle hits the S4 counter, see Sec. 3,
- (v) storage of the simulated events in a file which has the same format as the raw data,
- (vi) reconstruction of the simulated events with the same reconstruction chain as used for the real data and
- (vii) matching of reconstructed with simulated tracks based on the cluster positions.

It should be underlined that only inelastic p+p interactions in the hydrogen within the target cell were simulated and recon-

structed. Thus the simulation based corrections can be applied only for inelastic events. The contribution of elastic events is removed by the event selection cuts (see Sec. 5.1).

5 Data selection and analysis

This section presents the procedures used for data analysis [23]. These consist of the following steps: applying event and particle selections, obtaining uncorrected experimental results, evaluation of correction factors based on simulations, and finally calculation of the corrected results.

These steps are described in the subsections below.

5.1 Event and particle selection

5.1.1 Event selection criteria

The events selected for the analysis had to satisfy the conditions:

- (i) event is recorded with the interaction trigger,
- (ii) the beam particle trajectory is measured in at least one of BPD-1 or BPD-2 and in the BPD-3 detector,
- (iii) no off-time beam particle is detected within $\pm 1 \mu\text{s}$ around the trigger particle,
- (iv) there is at least one track reconstructed in the TPCs and fitted to the interaction vertex,
- (v) the vertex z position (fitted using the beam and TPC tracks) is not farther away than 10 cm from the centre of the LHT,
- (vi) events with a positively charged track with absolute momentum (p) close to the beam momentum ($p > p_{\text{beam}} - 3 \text{ GeV}/c$) are rejected to exclude elastic interactions.

The data statistics for analysed reactions is shown in Table 1.

	No cuts	Cuts applied
20 GeV/c	1320 k	176 k (13%)
31 GeV/c	3134 k	756 k (24%)
40 GeV/c	5238 k	1444 k (28%)
80 GeV/c	4500 k	1343 k (30%)
158 GeV/c	3537 k	1373 k (39%)

Table 1 Event statistics before and after event cuts for p+p data. See text for detailed description of the cuts.

5.1.2 Track selection criteria and acceptance

The tracks selected for the analysis had to satisfy the conditions:

- (i) track fit converged,
- (ii) the total number of reconstructed points on the track should be at least 30,
- (iii) the sum of the number of reconstructed points in VTPC-1 and VTPC-2 should be at least 15 or the number of reconstructed points in the GAP-TPC should be at least 5,
- (iv) the distance between the track extrapolated to the interaction plane and the interaction point (impact parameter) should be smaller than 4 cm in the horizontal (bending) plane and 2 cm in the vertical (drift) plane,
- (v) transverse momentum of particle track should be lower than 1.5 GeV/c,
- (vi) tracks with dE/dx and total momentum values characteristic for electrons are rejected.

Model simulations were performed in 4π acceptance, thus the NA61/SHINE detector acceptance filter needs to be applied before comparisons with data. The detector acceptance was defined as a three-dimensional matrix (p, p_T, ϕ) of 1 and 0 depending on whether the bin was or was not populated by particles reconstructed and accepted in the events (see Ref. [14]).

5.2 Raw correlation function

Uncorrected two-particle correlation functions were evaluated using Eq. 2 for events and tracks selected according to the criteria stated above. This was done by calculating differences between each accepted track in pseudorapidity and azimuthal angle with every other track in the same event. The differences were evaluated in the centre-of-mass (CMS) frame and accumulated in 2-dimensional bins of $(\Delta\eta, \Delta\phi)$. To increase statistics, the $\Delta\phi$ range was folded, i.e. for $\Delta\phi$ larger than π its value was recalculated as $2\pi - \Delta\phi$. This is allowed due to the symmetry of the p+p reactions. The transformation from laboratory (LAB) to CMS frame was performed assuming the pion mass for all produced particles.

The uncorrelated reference was constructed by mixing particles between events with two main constraints: a) the multiplicity distribution of mixed data had to be exactly the same as the original; b) mixed events could not contain two particles from the same

original (data) event. Example distributions of number of pairs of charged particles in bins of $(\Delta\eta, \Delta\phi)$ before and after mixing are shown in Fig. 2. The structures in the mixed events distributions are due to detector acceptance.

Examples of uncorrected correlation functions for all charge pairs² are presented in the left column of Fig. 3.

5.3 Corrections and uncertainties

5.3.1 Correction for reconstruction inefficiency and trigger-bias

In order to correct the results for biases due to trigger and off-line event and track selection, detection efficiency, contribution of weak decays and secondary interaction products, the same analysis was also performed on simulated data. Correction factors $\text{Corr}(\Delta\eta, \Delta\phi)$ were calculated bin-by-bin as the ratio of the correlation functions for simulated events from the EPOS [20] model (“pure”) and the same events after processing through GEANT [22] detector simulation and reconstruction (“rec”), and filtered using event and track selection cuts.

$$\text{Corr}(\Delta\eta, \Delta\phi) = \frac{C_{\text{pure}}^{\text{sim}}(\Delta\eta, \Delta\phi)}{C_{\text{rec}}^{\text{sim}}(\Delta\eta, \Delta\phi)}, \quad (3)$$

where $C_{\text{pure}}^{\text{sim}}$ and $C_{\text{rec}}^{\text{sim}}$ are the correlation function $C(\Delta\eta, \Delta\phi)$ obtained for simulated events before and after detector simulation and reconstruction. For both “pure” and “rec” events the NA61/SHINE acceptance filter was applied.

Corrected results $C(\Delta\eta, \Delta\phi)$ were obtained by multiplying the uncorrected correlation function by the corresponding correction, namely:

$$C(\Delta\eta, \Delta\phi) = C^{\text{raw}}(\Delta\eta, \Delta\phi) \cdot \text{Corr}(\Delta\eta, \Delta\phi). \quad (4)$$

Correction factors $\text{Corr}(\Delta\eta, \Delta\phi)$ and corrected correlation functions $C(\Delta\eta, \Delta\phi)$ for all charge pairs in p+p interactions at 20 and 158 GeV/c are shown in Fig. 3 as examples. Their values range between 0.9 and 1.1.

5.3.2 Statistical uncertainties

Statistical uncertainties of the correlation function are calculated as:

$$\sigma^2(C) = \sqrt{[\text{Corr} \cdot \sigma(C^{\text{raw}})]^2 + [C^{\text{raw}} \cdot \sigma(\text{Corr})]^2} \quad (5)$$

in each $(\Delta\eta, \Delta\phi)$ bin. Statistical uncertainties are found to be approximately independent of $\Delta\phi$ but increase significantly with increasing $\Delta\eta$. Depending on the beam momentum and charge

²Pairs of all, unlike-sign, positively, negatively charged particles are denoted by all, unlike-sign (+−), positive (++) , negative (−−) charge pairs for brief.




Fig. 2 Example distributions of number of pairs of all charge pairs in $(\Delta\eta, \Delta\phi)$ from the data (left) and mixed (right) events in p+p interactions in 158 GeV/c.




Fig. 3 Examples of uncorrected correlation functions from data (left plots), correction factors (centre plots) and corrected two-particle correlation functions $C(\Delta\eta, \Delta\phi)$ (right plots) for all charge pairs. Results are for p+p interactions at 20 GeV/c (top) and 158 GeV/c (bottom).

combination the statistical uncertainties of the C function in individual $(\Delta\eta, \Delta\phi)$ bins are about 5% for $\Delta\eta \approx 0$ (all charge pairs at top beam momentum) to over than 20% for $\Delta\eta \approx 3$ (negatively charge pairs at the lowest beam momentum).

5.3.3 Estimation of systematic uncertainties

In order to estimate systematic uncertainties the data were analysed with loose and tight event and track selection cuts. The standard set of cuts was presented in Sec. 5.1 and is tabulated in Table 2 together with loose and tight cuts.

Results for both sets of cuts were subtracted bin-by-bin (loose – tight). Since the differences in all bins follow Gaussian distributions with mean close to 0, the systematic uncertainties were esti-

mated as the standard deviation of the distribution. The procedure was performed for all charge combinations (all charge, unlike-sign, positively and negatively charge pairs) and at 20, 40 and 158 GeV/c. The systematic uncertainties are generally below 1%. The exception are correlation functions at 20 GeV/c at higher $\Delta\eta$, where the systematic uncertainties are about 5%.

6 Results and discussion

6.1 Two-particle correlation function $C(\Delta\eta, \Delta\phi)$

The corrected correlation functions for all charge pair combinations (all charge pairs, unlike-sign pairs, positively and negatively

Event cuts			
	Loose	Standard	Tight
Event with interaction trigger		applied	
BPD		applied	
No off-time beam particles	disabled	$< \pm 1 \mu\text{s}$	$< \pm 5 \mu\text{s}$
At least one track in TPCs		applied	
Vertex position z	$\pm 11 \text{ cm}$	$\pm 10 \text{ cm}$	$\pm 7 \text{ cm}$
Elastic event		applied	
Track cuts			
Charge $\neq 0$		applied	
Total TPC points	≥ 10	≥ 30	
VTPC (GTPC) points	$> 10(5)$	$\geq 15(5)$	$\geq 30(6)$
$ B_x $	$\leq 5 \text{ cm}$	$\leq 4 \text{ cm}$	$\leq 1 \text{ cm}$
$ B_y $	$\leq 2.5 \text{ cm}$	$\leq 2 \text{ cm}$	$\leq 0.5 \text{ cm}$
p_T cut		applied	
e^-e^+ cut		applied	

Table 2 Event (top) and track (bottom) selection cuts. The standard cuts (centre) are used to obtain the final results, whereas the loose (left) and tight (right) cuts are employed to estimate systematic uncertainties (see Sec. 5.1.1 and 5.1.2, respectively).

charge pairs) are presented in Figs. 4, 5, 6 and 7, respectively. Their values lie in the range between 0.8 and 1.4. Vanishing two-particle correlations result in $C = 1$.

Statistical and systematic uncertainties were calculated using the procedures described in Secs. 5.3.2 and 5.3.3. The numerical values of the correlation functions and of statistical and systematic uncertainties are available in Ref. [24].

The main features of the results are:

- (i) A maximum at $(\Delta\eta, \Delta\phi) \approx (0, \pi)$, most prominent for all charge and unlike-sign pairs and significantly weaker for like-sign pairs. The most probable explanation is the contribution from resonance decays, mostly from abundantly produced $\rho^0 \rightarrow \pi^+ + \pi^-$. The weaker maximum for positive charge pairs can be attributed to e.g. Δ^{++} resonance decay. No such maximum is observed for negative charge pairs consistent with the fact that there are almost no resonances decaying into two or more negative charge pairs.
- (ii) A $\cos(\Delta\phi)$ modulation appearing as a minimum near $\Delta\phi = 0$ and maximum near $\Delta\phi = \pi$ at all values of $\Delta\eta$ for all combinations of charges. Stronger in all charge and unlike-sign pairs, weaker but still visible in positive charge and barely noticeable for negative charge pairs. The structure is probably due to momentum conservation.
- (iii) A Gaussian-like enhancement around $\Delta\eta = 0$ along the full $\Delta\phi$ range, clearly visible for all charge and unlike-sign pairs, however significantly weaker, but still noticeable in like-sign pairs. This feature may be connected with string fragmentation or flux-tube fragmentation [25].
- (iv) A hill around $(\Delta\eta, \Delta\phi) = (0, 0)$ for like-sign pairs. For positive charged pairs it grows with increasing beam momen-

tum, but it is independent of beam momentum for negative charge pairs. Since the products of γ decay were rejected during the analysis, the hill is probably caused by Bose-Einstein statistics.

- (v) Clearly, there is no jet-like peak at $(\Delta\eta, \Delta\phi) = (0, 0)$. This indicates that hard scattering effects are absent or below the background threshold at SPS energies.

6.2 Projections onto $C(\Delta\eta)$ and $C(\Delta\phi)$

To make the results easier to compare quantitatively, slices of the two-dimensional correlation function are also presented. The $C(\Delta\eta)$ function was obtained by projecting $C(\Delta\eta, \Delta\phi)$ onto $\Delta\eta$ in the $\Delta\phi$ intervals: $0 < \Delta\phi < \pi/4$ and $3\pi/4 < \Delta\phi < \pi$. The results for 20, 40, and 158 GeV/c beam momenta and all charge, unlike-sign and positive charged pairs, together with uncertainties as well as predictions of the EPOS and UrQMD models are presented in Fig. 8. Note that due to the projection the statistical uncertainties of the projected correlation function are considerably smaller than for individual two-dimensional bins.

The two-particle correlation function $C(\Delta\phi)$ was obtained from projection of $C(\Delta\eta, \Delta\phi)$ onto the $\Delta\phi$ axis in two $\Delta\eta$ intervals: $0 < \Delta\eta < 1$ and $2 < \Delta\eta < 3$. The results are presented in Fig. 9. The tendencies shown in Figs. 8 and 9 are reproduced well by the EPOS model.

6.3 Acceptance effects

Acceptance effects were studied using events generated with the EPOS model. Namely, a mixed (uncorrelated) background distribution $M(\Delta\eta, \Delta\phi)$ was produced for EPOS events in full and the NA61/SHINE detector acceptance. Figure 10 shows the comparison between mixed events for 4π acceptance (left) and NA61/SHINE acceptance (centre) for 20 and 158 GeV/c. The right column of Fig. 10 displays the ratio of these two distributions. One finds large effects of the limited NA61/SHINE acceptance which are strongest at 20 GeV/c. It is therefore essential to apply the NA61/SHINE acceptance filter to model calculations before comparing to the measurements.

7 Comparison with models

7.1 Comparison with the EPOS and UrQMD models

In order to compare the experimental results and predictions of the EPOS 1.99 [20] and UrQMD 3.4 [26, 27] models the ratio of correlation functions obtained from data and models was calculated. The same acceptance was used for the NA61/SHINE data and for the EPOS and UrQMD models (see Ref. [14]). The results for unlike-sign and positive charge pairs are shown in Fig. 11. In general, EPOS 1.99 reproduces the experimental results from p+p




Fig. 4 Two-particle correlation function $C(\Delta\eta, \Delta\phi)$ for all charge pairs in inelastic p+p interactions at 20-158 GeV/c. The correlation function is mirrored around $(\Delta\eta, \Delta\phi) = (0, 0)$.




Fig. 5 Two-particle correlation function $C(\Delta\eta, \Delta\phi)$ for unlike-sign pairs in inelastic p+p interactions at 20-158 GeV/c. The correlation function is mirrored around $(\Delta\eta, \Delta\phi) = (0, 0)$.




Fig. 6 Two-particle correlation function $C(\Delta\eta, \Delta\phi)$ for positive charge pairs in inelastic p+p interactions at 20-158 GeV/c. The correlation function is mirrored around $(\Delta\eta, \Delta\phi) = (0, 0)$.




Fig. 7 Two-particle correlation function $C(\Delta\eta, \Delta\phi)$ for negative charge pairs in inelastic p+p interactions at 20-158 GeV/c. The correlation function is mirrored around $(\Delta\eta, \Delta\phi) = (0, 0)$.




Fig. 8 Two-particle correlation function $C(\Delta\eta)$ obtained from projection of $C(\Delta\eta, \Delta\phi)$ onto the $\Delta\eta$ axis for two subranges of $\Delta\phi$. Left column shows the results for all charge pairs, middle column – unlike-sign pairs, right column – positive charge pairs. Vertical bars denote statistical and shaded regions denote systematic uncertainties. Predictions of the EPOS model are shown by solid curves and the UrQMD model by dotted curves.

interactions well (see first and third column in Fig. 11). However, the model does not reproduce the peak of about 20% at $(\Delta\eta, \Delta\phi) = (0, 0)$ seen for positive charge pairs (third column in Fig. 11) at 40, 80 and 158 GeV/c. This demonstrates an excess of the number of real data over model pairs in that region. Note, that the EPOS model does not include either Bose-Einstein

correlations or Coulomb interactions which are expected to produce this kind of correlation. Moreover a weak enhancement of the ratio of about 10% is seen near $\Delta\eta \approx 0$ and $\Delta\phi \approx \pi$ and $\frac{3}{4}\pi$ (first column in Fig. 11). This suggests a slight underestimate of the correlation in the valleys of the $\cos(\Delta\phi)$ modulation in EPOS.




Fig. 9 Two-particle correlation function $C(\Delta\phi)$ obtained from projection of $C(\Delta\eta, \Delta\phi)$ onto the $\Delta\phi$ axis for two subranges of $\Delta\eta$. Left column shows the results for all charge pairs, middle column – unlike-sign pairs, right column – positive charge pairs. Vertical bars denote statistical and shaded regions denote systematic uncertainties. Predictions of the EPOS model are shown by solid curves and the UrQMD model by dotted curves.

The ratio of the experimental results and the predictions of the UrQMD 3.4 model is shown in the second and fourth column of Fig. 11. The UrQMD model reproduces data less well than EPOS. The most visible discrepancy is a $\Delta\eta$ -wide suppression of the ratio in unlike-sign pairs (second column in Fig. 11) which suggests that UrQMD generates 25 - 30% stronger long-range near-side

($\Delta\phi \approx 0$) correlations than existing in the data. Another inconsistency between data and model results can be seen in positive charge correlations (fourth column in Fig. 11) where a suppression of up to 25% is present on the away-side ($\Delta\phi \approx \pi$), strongest at the three lowest beam momenta, as well as an enhancement of more than 10% at large $\Delta\eta$ on the same side in azimuthal angle.




Fig. 10 NA61/SHINE detector acceptance effects studied with the EPOS [20] model. Left plots show mixed event distributions $M(\Delta\eta, \Delta\phi)$ for 4π acceptance (left) and for NA61/SHINE acceptance (centre). Right plots show the ratio of correlation functions calculated with both acceptances. Top row shows results for 20 GeV/c, bottom row for 158 GeV/c.

8 Summary

Two-particle correlations of charged particles in azimuthal angle and pseudorapidity were measured in inelastic p+p collisions at 20 - 158 GeV/c by the NA61/SHINE experiment at the CERN SPS. The results show structures which can be connected to phenomena such as resonance decays, momentum conservation and Bose-Einstein correlations. A comparison with the EPOS and UrQMD models was performed. The predictions of the EPOS model are close to the NA61/SHINE data whereas the results from UrQMD show larger discrepancies.

Acknowledgements We would like to thank the CERN EP, BE and EN Departments for the strong support of NA61/SHINE.

This work was supported by the Hungarian Scientific Research Fund (grants OTKA 68506 and 71989), the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, the Polish Ministry of Science and Higher Education (grants 667/N-CERN/2010/0, NN 202 48 4339 and NN 202 23 1837), the Polish National Center for Science (grants 2011/03/N/ST2/03691, 2013/11/N/ST2/03879, 2014/13/N/ST2/02565, 2014/14/E/ST2/00018, 2015/18/M/ST2/00125, and 2015/19/N/ST2/01689), the Foundation for Polish Science — MPD program, co-financed by the European Union within the European Regional Development Fund, the Federal Agency of Education of the Ministry of Education and Science of the Russian Federation (SPbSU research grant 11.38.242.2015), the Russian Academy of Science and the Russian Foundation for Basic Research (grants 08-02-00018, 09-02-00664 and 12-02-91503-CERN), the Ministry of Education, Culture, Sports, Science and Technology, Japan, Grant-in-Aid for Scientific Research (grants 18071005, 19034011, 19740162, 20740160 and 20039012), the German Research Foundation (grant GA 1480/2-2), the EU-funded Marie Curie Outgoing Fellowship, Grant PIOF-GA-2013-624803, the Bulgarian Nuclear Regu-

latory Agency and the Joint Institute for Nuclear Research, Dubna (bilateral contract No. 4418-1-15/17), Ministry of Education and Science of the Republic of Serbia (grant OI171002), Swiss Nationalfonds Foundation (grant 200020117913/1), ETH Research Grant TH-01 07-3 and the U.S. Department of Energy.

References

1. <https://shine.web.cern.ch/>.
2. N. Abgrall et al. NA61/SHINE facility at the CERN SPS: beams and detector system. *JINST*, 9:P06005, 2014.
3. M. Gazdzicki and P. Seyboth. Search for Critical Behaviour of Strongly Interacting Matter at the CERN Super Proton Synchrotron. *Acta Phys. Polon.*, B47:1201, 2016.
4. K. Eggert et al. Angular Correlations Between the Charged Particles Produced in pp Collisions at ISR Energies. *Nucl.Phys.*, B86:201, 1975.
5. B. Alver et al. Cluster properties from two-particle angular correlations in p + p collisions at $s^{*(1/2)} = 200$ -GeV and 410-GeV. *Phys.Rev.*, C75:054913, 2007.
6. R.J. Porter and T.A. Trainor. Correlation structures from soft and semi-hard components in p - p collisions at $s^{*(1/2)} = 200$ -GeV. *Acta Phys.Polon.*, B36:353–359, 2005.
7. R. J. Porter and T. A. Trainor. Soft and hard components of two-particle distributions on $(y(t), \eta(t), \phi)$ from p-p collisions at $s^{*(1/2)} = 200$ -GeV. In *Ultra-relativistic nucleus-nucleus collisions. Proceedings, 17th International Confer-*




Fig. 11 The ratio of correlation functions $C(\Delta\eta, \Delta\phi)$ for data and models. The two left columns show results for unlike-sign pairs and the two right columns results for positive charge pairs. Each row shows results for one beam momentum.

- ence, *Quark Matter 2004, Oakland, USA, January 11-17, 2004*, 2004.
8. B. Abelev et al. Multiplicity dependence of jet-like two-particle correlations in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. *Phys.Lett.*, B741:38–50, 2015.
 9. B. Alver et al. System size dependence of cluster properties from two-particle angular correlations in Cu+Cu and Au+Au collisions at $s(NN)^{1/2} = 200$ -GeV. *Phys.Rev.*, C81:024904, 2010.
 10. G. Agakishiev et al. Anomalous centrality evolution of two-particle angular correlations from Au-Au collisions at $\sqrt{s_{NN}} = 62$ and 200 GeV. *Phys.Rev.*, C86:064902, 2012.
 11. Vardan Khachatryan et al. Observation of Long-Range Near-Side Angular Correlations in Proton-Proton Collisions at the LHC. *JHEP*, 1009:091, 2010.
 12. Vardan Khachatryan et al. Measurement of long-range near-side two-particle angular correlations in pp collisions at $\sqrt{s} = 13$ TeV. *Phys. Rev. Lett.*, 116(17):172302, 2016.
 13. Serguei Chatrchyan et al. Observation of long-range near-side angular correlations in proton-lead collisions at the LHC. *Phys.Lett.*, B718:795–814, 2013.
 14. <https://edms.cern.ch/document/1700774/1>.
 15. Georges Aad et al. Measurement of inclusive two-particle angular correlations in pp collisions with the ATLAS detector at the LHC. *JHEP*, 1205:157, 2012.
 16. Malgorzata Janik. $\Delta\eta\Delta\phi$ angular correlations in pp collisions at the LHC registered by the ALICE experiment. *PoS, WPCF2011:026*, 2011.
 17. C. Bovet et al. The Cedar Counters for Particle Identification in the SPS Secondary Beams: A Description and an Operation Manual. 1982.
 18. C. Strabel. Evaluation of Particle Yields in 30 GeV Proton-Carbon Inelastic Interactions for Estimating the T2K Neutrino Flux. *ETH-19538, CERN-THESIS-2011-295*, 2011.
 19. N. Abgrall. Report from the NA61/SHINE experiment at the CERN SPS. Technical Report CERN-SPSC-2014-031. SPSC-SR-145, CERN, Geneva, Oct 2014.
 20. K. Werner, F.-M. Liu, and T. Pierog. Parton ladder splitting and the rapidity dependence of transverse momentum spectra in deuteron-gold collisions at RHIC. *Phys.Rev.*, C74:044902, 2006.
 21. N. Abgrall, 2011. PhD. Thesis, University of Geneva, Switzerland, <https://edms.cern.ch/document/1172455/1>.
 22. R. Brun and F. Carminati. Geant detector description and simulation tool, cern program library long writeup w5013, 1993. <http://wwwasdoc.web.cern.ch/wwwasdoc/geant/geantall.html>.
 23. B. Maksiak. *Azimuthal angle and pseudorapidity correlations in p+p and Pb+Pb collisions at energies of SPS accelerator (in preparation)*. PhD thesis, Faculty of Physics, Warsaw University of Technology, 2016.
 24. Numerical values of NA61/SHINE two-particle correlations analysis <https://edms.cern.ch/document/1701318/1>.
 25. C.-Y. Wong. Signature of the Fragmentation of a Color Flux Tube. *Phys. Rev.*, D92:074007, 2015.
 26. S.A. Bass et al. Microscopic models for ultrarelativistic heavy ion collisions. *Prog.Part.Nucl.Phys.*, 41:255–369, 1998.
 27. M. Bleicher et al. Relativistic hadron hadron collisions in the ultrarelativistic quantum molecular dynamics model. *J.Phys.*, G25:1859–1896, 1999.